

**ART\_RS field**

**Table of Contents**

<b>ART_RS field</b>	<b>1</b>
Coding Table	2

containing table

[tblART](#)

explanation of variable

Reason for stopping treatment

format of data

character. see [coding table](#) for valid codings.

exists since HICDEP version

[1.30](#)

## Coding Table

[Download this table as CSV file](#) [Download this table as XML file](#)

Code	Reason for Stopping Treatment
1	Treatment failure (i.e. virological, immunological, and/or clinical failure)
1.1	Virological failure
1.2	Partial virological failure
1.3	Immunological failure - CD4 drop
1.4	Clinical progression
2	Abnormal fat redistribution
3	Concern of cardiovascular disease
3.1	Dyslipidaemia
3.2	Cardiovascular disease
4	Hypersensitivity reaction
5	Toxicity, predominantly from abdomen/G-I tract
5.1	Toxicity - GI tract
5.2	Toxicity - Liver
5.3	Toxicity - Pancreas
6	Toxicity, predominantly from nervous system
7	Toxicity, predominantly from kidneys
8	Toxicity, predominantly from endocrine system
8.1	Diabetes
9	Haematological toxicity (anemia etc.)
10	Hyperlactataemia/lactic acidosis
88	Death
90	Side effects - any of the above but unspecified
90.1	Comorbidity
91	Toxicity, not mentioned above
92	Availability of more effective treatment (not specifically failure or side effect related)
92.1	Simplified treatment available
92.2	Treatment too complex
92.3	Drug interaction
93	Structured Treatment Interruption (STI)
93.1	Structured Treatment Interruption (STI) - at high CD4

94	Patient's wish/decision, not specified above
94.1	Non-compliance
95	Physician's decision, not specified above
96	Pregnancy
97	Study treatment
98	Other causes, not specified above
99	Unknown

Although most reasons mentioned above could be coded in ICD-10 codes, it is still certain that many reasons will never be defined in terms of ICD-10 (STI, physicians and patients decision), so to avoid a mixture of two different coding systems it is advised to go with the above list and build upon that for now.